

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/IT2006/000049

International filing date: 27 January 2006 (27.01.2006)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/647,498
Filing date: 27 January 2005 (27.01.2005)

Date of receipt at the International Bureau: 25 April 2006 (25.04.2006)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

Paolo Jorio
Roberto Prato
Luigi Boggio
Rinaldo Plebani
Corrado Modugno
Simone Bongiovanni
Luigi Franzolin
Elena Cerbaro
M. Cristina Baldini
Giancarlo Revelli
Mauro Eccetto
Raffaele Borrelli
Fabio D'Angelo
Mirko Bergadano
Matteo Bellemo
Matteo Maccagnan
Daniele Cernuzzi
Stefano Manconi Dirrigl
Raffaella Barbuto
Robert Scotti (US IP Att.)
Satsuki Takasaka
Francesco Fiussello
Giovanni Lo Cigno

Andrea Bernotti
Paolo Lovino
Cesare Bosman
Stefania Berta
Bruno Zamprognò
Simone Mangini

Edoardo Mola
Lorenzo Nannucci
Manuela Giannini
Cristina Rolando
Andrea Fiorini
Michele Di Sciuvia
Lidia Casciano
Alessandro Putzolu
Erika Andreotti
Gerardo Valeriano

Legal Advisers:
Claudio Costa
Maria Teresa Saguatti
Chiara Luzzato
Alberto Improda
Raffaella Arista



STUDIO TORTA

JORIO, PRATO, BOGGIO & Partners

Via Viotti, 9 - 10121 Torino - Italy
Tel. +39-011.561.13.20; Fax +39-011.562.21.02
Videoconference +39-011.518.42.48 (6 ISDN channels H.320)
E-mail: info@studiotorta.it Internet: www.studiotorta.com

Industrial Property Consultants - Established in 1879
European and Italian Patent Attorneys
European and Italian Trade Mark and Design Attorneys
TURIN MILAN BOLOGNA TREVISO RIMINI ROME

**International Bureau of WIPO
PCT Receiving Office Section**
34, Chemin des Colombettes
CH-1211 GENEVA 20
SWITZERLAND
(Svizzera)

Torino, April 18, 2006

=====

Our case E4031/06-WO-cob

Dear Sirs,

RE: International Application No. PCT/IT2006/000049 filed on January 27, 2006
in the name of **ALMA MATER STUDIORUM - UNIVERSITA' DI BOLOGNA**

With reference to the application in object, we enclose herewith the priority document of the basic US Patent application.

Yours faithfully,

Dr. Ing. Elena CERBARO


10121 Torino
Via Viotti, 9
Tel. +39-011.561.13.20
Fax. +39-011.562.21.02

20123 Milano
Corso Magenta, 56
Tel. +39-02.48.01.42.16
Fax +39-02.48.01.50.82

40133 Bologna
Via Emilia Ponente, 34
Tel. +39-051.38.91.22
Fax +39-051.38.90.30

31100 Treviso
Viale Appiani, 26
Tel. +39-0422.22.199
Fax +39-0422.23.316

00187 Roma
Via Due Macelli, 47
Tel. +39-06.67.91.589
Fax +39-06.67.97.747



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

February 09, 2006


THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/647,498

FILING DATE: January 27, 2005

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS US60/647,498

By Authority of the
Under Secretary of Commerce for Intellectual Property
and Director of the United States Patent and Trademark Office



P. SWAIN
Certifying Officer

18379 U.S. PTO

PTO/SB/16 (12-04)

Approved for use through 07/31/2006. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. EV333073539US

12900 U.S. PTO
60/647498



INVENTOR(S)		
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)
Michela	ROSINI	Bologna, Italy
Additional inventors are being named on the <u>1</u> separately numbered sheets attached hereto		
TITLE OF THE INVENTION (500 characters max):		
MULTIPOTENT ANTI-ALZHEIMER DRUGS		
CORRESPONDENCE ADDRESS		
Direct all correspondence to:		
<input type="checkbox"/> The address corresponding to Customer Number: 		
OR		
<input checked="" type="checkbox"/> Firm or Individual Name <u>Law Offices of Albert Wai-Kit Chan, LLC</u>		
Address <u>World Plaza, Suite 604</u> <u>141-07 20th Avenue</u>		
City <u>Whitestone</u>	State <u>New York</u>	Zip <u>11357</u>
Country <u>U.S.A.</u>	Telephone <u>(718)-357-8836</u>	Fax <u>(718)-357-8615</u>
ENCLOSED APPLICATION PARTS (check all that apply)		
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76		
<input checked="" type="checkbox"/> Specification Number of Pages <u>19</u>		
<input checked="" type="checkbox"/> Drawing(s) Number of Sheets <u>4</u>		
<input type="checkbox"/> CD(s), Number of CDs _____		
<input checked="" type="checkbox"/> Other (specify) <u>Express Mail Certificate of Mailing bearing Label No. EV333073539US</u>		
Application Size Fee: If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$250 (\$125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).		
METHOD OF PAYMENT OF FILING FEES AND APPLICATION SIZE FEE FOR THIS PROVISIONAL APPLICATION FOR PATENT		
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.		TOTAL FEE AMOUNT (\$)
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the filing fee and application size fee (if applicable).		<u>200.00</u>
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached		
<input checked="" type="checkbox"/> The Director is hereby authorized to charge the filing fee and application size fee (if applicable) or credit any overpayment to Deposit Account Number: <u>50-1891</u> . A duplicative copy of this form is enclosed for fee processing.		
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.		
<input type="checkbox"/> No.		
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____		

SIGNATURE Albert Wai Kit Chan

Date 1/27/05

TYPED or PRINTED NAME Albert Wai-Kit Chan

REGISTRATION NO. 36,479
(if appropriate)

TELEPHONE (718) 357-8836

Docket Number: 1033-PRO

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

PROVISIONAL APPLICATION COVER SHEET
Additional Page

PTO/SB/16 (12-04)

Approved for use through 07/31/2006. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

First Named Inventor	Michela ROSINI	Docket Number	1033-PRO
----------------------	----------------	---------------	----------

INVENTOR(S)/APPLICANT(S)

INVENTOR(S)/APPLICANT(S)		
Given Name (first and middle [if any])	Family or Surname	Residence (City and either State or Foreign Country)
Vincenza	ANDRISANO	Bologna, Italy
Manuela	BARTOLINI	Bologna, Italy
Carlo	MELCHIORRE	Bologna, Italy

Number 2 of 2

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s) : ROSINI, et al.
U.S. Serial No.: Not Yet Known
Filed : January 27, 2005
For : MULTIPOTENT ANTI-ALZHEIMER DRUGS
Law Offices of Albert Wai-Kit Chan, LLC
World Plaza, Suite 604
141-07 20th Avenue
Whitestone, NY 11357
January 27, 2005

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450
Mail Stop Provisional Application

Sir:


EXPRESS MAIL CERTIFICATE OF MAILING
IN CONNECTION WITH THE ABOVE-IDENTIFIED APPLICATION

"Express Mail" mailing label number: EV333073539US

Date of Deposit: January 27, 2005

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. §1.10 on the date indicated above and is addressed to Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450, Mail Stop Provisional Application.


Printed Name: Alex Chay Chang

Respectfully submitted,

Albert Wai-Kit Chan
Registration No. 36,479
Attorney for Applicant(s)
Law Offices of
Albert Wai-Kit Chan, LLC
World Plaza, Suite 604
141-07 20th Avenue
Whitestone, New York 11357
Tel: (718) 357-8836
Fax: (718) 357-8615
e-mail: kitchenlaw@aol.com

*Application
for
United States Letters Patent*

To all whom it may concern:

Be it known that Michela ROSINI, Vincenza ANDRISANO, Manuela BARTOLINI, and Carlo MELCHIORRE

have invented certain new and useful improvements in

MULTIPOTENT ANTI-ALZHEIMER DRUGS

of which the following is a full, clear, and exact description.

Multipotent Anti-Alzheimer Drugs

Abstract:

The coupling of two different pharmacophores, each endowed with different biological properties, afforded the hybrid compound lipocrine (7), whose biological profile was markedly improved relative to those of prototypes tacrine and lipoic acid. Lipocrine is the first compound that inhibits the catalytic activity of AChE and AChE-induced amyloid- β aggregation and protects against reactive oxygen species. Thus, it emerged as a valuable pharmacological tool to investigate Alzheimer's disease and as a promising lead compound for new anti-Alzheimer drugs.

~~~~~

Alzheimer's disease (AD), the most common cause of dementia, is a complex neurological affection that is clinically characterized by loss of memory and progressive deficits in different cognitive domains. The consistent neuropathologic hallmark of the disorder, generally noted on postmortem brain examination, is a massive deposit of aggregated protein breakdown products, amyloid- $\beta$  (A $\beta$ ) plaques and neurofibrillary tangles. Even if the primary cause of AD is still speculative, A $\beta$  aggregates are thought to be mainly responsible for the devastating clinical effects of the disease.<sup>1</sup> In recent years, significant research has been devoted to the role of free radical formation, oxidative cell damage, and inflammation in the pathogenesis of AD, providing new promising targets and validated animal models.<sup>2</sup> To date, however, the enhancement of the central cholinergic function is the only clinically effective approach.<sup>3,4</sup> The intensive research of drugs that can improve the cholinergic transmission in AD has produced so far four approved acetylcholinesterase (AChE) inhibitors, that is, tacrine (TC),<sup>5</sup> donepezil,<sup>6</sup> rivastigmine,<sup>7</sup> and galantamine.<sup>8</sup> However, these drugs have been approved for the symptomatic treatment of AD because they do not address the etiology of the disease for which they are used.


It is therefore necessary to discover pharmacological instruments that are able to act as far upstream as possible in the neurodegenerative cascade and, because of the multifaceted etiology of AD, able to hit different selected targets. The aim of this communication is to provide new multipotent compounds, i.e., single molecules that can exhibit more pharmacological properties simultaneously,



such as the enhancement of the cholinergic transmission and inhibition of A $\beta$  accumulation and oxidative stress, leading to a synergic and effective treatment of AD.


To this end, we applied a design strategy in which distinct pharmacophores of two different drugs were combined in the same structure leading to hybrid molecules. In principle, each pharmacophore of these new drugs should retain the ability to interact with its specific site(s) on the target and consequently to produce specific pharmacological responses that taken together should block or hopefully cure the neurodegenerative process leading to AD. To obtain proof of concept for this proposal, we chose TC and lipoic acid (LA) as prototype drugs to be combined in the same structure because of their well-established biological properties (Figure 1). In fact, LA is a universal antioxidant,<sup>9-12</sup> which was shown to protect neurons against cytotoxicity induced by A $\beta$ <sup>13</sup> and to stabilize cognitive functions in patients with AD,<sup>14</sup> whereas TC was the first AChE inhibitor approved for AD treatment.<sup>5</sup> AChE is the enzyme involved in the hydrolysis of the neurotransmitter acetylcholine (ACh) at cholinergic synapses in the central and peripheral nervous system. Inhibitors of AChE activity promote an increase in the concentration and the duration of action of synaptic ACh, thus causing an enhancement of the cholinergic transmission through activation of the synaptic nicotinic and muscarinic receptors. However, achievement of potent inhibitors of the AChE catalytic site would not represent a significant improvement unless there is a concomitant inhibition of the peripheral anionic site (PAS) of the enzyme, which is associated with the neurotoxic cascade of AD through AChE-induced A $\beta$  aggregation.<sup>15,16</sup> For this reason, the introduction onto the TC structure of a side chain, namely, an LA fragment following the two routes shown in Figure 1, should hopefully combine antioxidant properties with the ability to interact with PAS.

The strategy has been successfully utilized for other compounds more than tacrine. The general structure of lipoic acid derivatives can be schematized as such:



Where n = 1-4;

Examples where R =:




The examples are compounds (1-8) reported in the enclosed document.

An other example is an inferior homologous of **8**:

**RM11**:  $n = 1$ ,  $R_1 = 3$ -ylmethylamine,  $R_2 = \text{NH}_2$ ,  $R_3 = 6$ -chloro;


b)



Example:

**MR106**:  $n = 4$ ;  $X = \text{CH}=\text{CH}$ ,  $R_1 = 2$ -MeO,  $m = 6$ ;  $R_2 = \text{Et}$ ,  $R_3 = \text{H}$ .

c)



$R_1 = \text{alkyl}$ ,  $R_2 = \text{alkyl}$ , alkylamine

$Y = \text{alkyl}$

$X = \text{OH}$ , O-carbamoyl, O-

An example is **MR112**:  $Y = \text{MeCH}$ ,  $R_1 = R_2 = \text{Me}$ ,  $X = \text{O-}$ .

Compounds **1-8** were synthesized by coupling tetrahydroacridine intermediates **9-16** with LA in the presence of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), according to Scheme 1 (see Supporting Information). No attempt was made to separate the enantiomers of **1-7** because it was reported that stereochemistry is not relevant for the protective effect of LA against oxidative cell damage.<sup>10</sup> In addition, concerning **8**, the diastereomers were not separated because it was the weakest AChE inhibitor among the investigated compounds.

To determine the potential interest of **1-8** for the treatment of AD, their AChE inhibitory activity was determined by the method of Ellman et al.<sup>17</sup> on human recombinant AChE. Moreover, to study further the biological profiles of **1-8**, their butyrylcholinesterase (BChE) inhibitory activity was also assessed by the same method on BChE from human serum. To allow comparison of the results, **15**, TC, and LA were used as the reference compounds (Table 1e). It is evident that all compounds were effective inhibitors of AChE and BChE and significantly more potent than prototype TC with the exception of **8**. Modifying the chain length between the two nitrogen atoms of the lateral chain,

affording 1-7, affected the affinity for AChE and BChE. Optimum inhibition of AChE was observed for 2, having three methylene units. As expected,<sup>18</sup> the insertion of a chlorine atom into the acrydine system, affording 7 (lipocrine), produced an 85-, 1676-, and 28-fold increase in AChE inhibition relative to 15, TC, and 2, respectively. As one would expect, LA did not inhibit either enzyme. The finding that 8 was 4308-fold less potent than 2 in inhibiting AChE activity suggests that the insertion of the lipoyl fragment on the nitrogen atom at position 3 of 16 resulted in a highly negative effect on the interaction mechanism with the enzyme.

Inhibition of AChE activity by 7 was very fast and not time-dependent because 50% of enzyme inactivation produced by 0.253 nM following a 1 min incubation was not significantly different ( $p > 0.01$ ) from the inhibition observed up to a 40 min incubation. The estimates of competitive inhibition constants  $K_i$  calculated for 7 and TC were  $0.155 \pm 0.046$  and  $0.151 \pm 0.016$   $\mu$ M, respectively, whereas the graphical analysis of steady-state inhibition data for 7 is shown in Figure 2. An analysis of the Lineweaver-Burk reciprocal plots of 7 reveals that there are an increasing slope and an increasing intercept with higher inhibitor concentration. The inhibitory behavior of 7, as deduced from Figure 2, is strictly similar to that displayed by some reported bis-tetrahydroaminoacridine inhibitors of AChE. These compounds bind simultaneously at the catalytic and the peripheral sites of AChE and are characterized by a linear mixed type of enzyme inhibition.<sup>19</sup> Therefore, we concluded that 7 causes a mixed type of inhibition, i.e., inhibition of both the active site and a second distal site of the enzyme. Once verified that 7 may interact also with PAS of AChE, we verified whether there is a concomitant inhibition of A $\beta$  aggregation induced by AChE through a thioflavin T-based fluorometric assay.<sup>16,20</sup> It turned out that TC and LA, i.e., the pharmacophoric moieties combined in 7, were not able to inhibit at 100  $\mu$ M the A $\beta$  aggregation induced by AChE, whereas 100  $\mu$ M 15 caused only a  $25 \pm 5\%$  inhibition. In contrast, 7 was only 3-fold less potent than propidium, which is the most potent inhibitor of AChE-induced A $\beta$  aggregation so far available,<sup>16</sup> as revealed by an analysis of their  $IC_{50}$  values of  $45.0 \pm 14.6$  and  $12.6 \pm 0.5$   $\mu$ M, respectively. Furthermore, 7 was significantly more potent than all the other AChE inhibitors ever tested,<sup>16</sup> including AP2238, an inhibitor purposely designed to bind the catalytic and the peripheral sites of AChE (35% of inhibition at 100  $\mu$ M).<sup>21</sup>

Clearly, this finding, together with the results observed for 15, TC, and LA separately, is relevant because an association of 100  $\mu$ M TC and 100  $\mu$ M LA or 100  $\mu$ M 15 and 100  $\mu$ M LA produced only a weak inhibition ( $15 \pm 6\%$  or  $30 \pm 7\%$ , respectively) of AChE-induced A $\beta$  aggregation, suggesting

that marked A $\beta$  aggregation inhibition may be achieved only when the two prototypes are combined into the same structure, as in 7.

The cytotoxicity effects of LA, 7, and 15 were first determined by colorimetric MTT [3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide] assay in human neuronal-like cells, SH-SY5Y, as described by Mosmann et al.<sup>22</sup> As reported in Figure 3, treatment of SH-SY5Y cells with LA and 7 (0.1-50  $\mu$ M) did not show modified cell viability. By contrast, the treatment of SH-SY5Y cells with 15 (0.1-50  $\mu$ M) produced a strong decrease of cell viability for 10  $\mu$ M (88%) and 50  $\mu$ M (99%). The intracellular antioxidant activity of LA, 7, and 15 against formation of reactive oxygen species (ROS) in SH-SY5Y cells after treatment with *tert*-butyl hydroperoxide, a compound used to induce oxidative damage, was then assessed. A range of concentrations of tested compounds that did not affect cell viability (0.1-50  $\mu$ M for LA and 7; 0.1-5  $\mu$ M for 15) were used. As shown in Table 2a, treatment of SH-SY5Y cells with LA showed a significant ( $p < 0.01$ ) decrease of ROS formation only with the highest concentration used (50  $\mu$ M), while the treatment with 7 produced a strong dose-dependent inhibitory effect on the ROS formation. Significant inhibitory effects by 7 with respect to basal values were reached for 5  $\mu$ M ( $p < 0.01$ ), 10  $\mu$ M ( $p < 0.001$ ), and 50  $\mu$ M ( $p < 0.001$ ). When treated with 15 (0.1-5  $\mu$ M), the neuronal cells did not show any difference on ROS formation. Taken together, these results show that LA and 7 did not affect the neuronal viability while 15 exerted neurotoxic effects. In addition, LA and 7 (but not 15) were able to protect neuronal cells against ROS formation evoked by oxidative stress, with 7 being the most active against ROS formation (64% inhibition at 50  $\mu$ M).

In conclusion, the present investigation has shown that it is possible to obtain multipotent drugs for the treatment of AD: lipocrine (7) emerged in in vitro models as an effective candidate to be investigated in vivo for its multiple biological properties, namely, inhibition of AChE and BChE activities, inhibition of AChE-induced A $\beta$  aggregation, and ability to protect cells against ROS.

## BIBLIOGRAPHY

1. Hardy, J.; Selkoe, D. J. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* **2002**, *297*, 353-356.
2. Capsoni, S.; Ugolini, G.; Comparini, A.; Ruberti, F.; Berardi, N.; Cattaneo, A. Alzheimer-like neurodegeneration in aged antinerve growth factor transgenic mice. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 6826-6831.

3. Bartus, R. T. On neurodegenerative diseases, models, and treatment strategies: lessons learned and lessons forgotten a generation following the cholinergic hypothesis. *Exp. Neurol.* **2000**, *163*, 495-529.
4. Terry, A. V., Jr.; Buccafusco, J. J. The cholinergic hypothesis of age and Alzheimer's disease-related cognitive deficits: recent challenges and their implications for novel drug development. *J. Pharmacol. Exp. Ther.* **2003**, *306*, 821-827.
5. Davis, K. L.; Powchik, P. Tacrine. *Lancet* **1995**, *345*, 625-630.
6. Bryson, H. M.; Benfield, P. Donepezil. *Drugs Aging* **1997**, *10*, 234-239, 240-241 (discussion).
7. Gabelli, C. Rivastigmine: an update on therapeutic efficacy in Alzheimer's disease and other conditions. *Curr. Med. Res. Opin.* **2003**, *19*, 69-82.
8. Sramek, J. J.; Frackiewicz, E. J.; Cutler, N. R. Review of the acetylcholinesterase inhibitor galanthamine. *Expert Opin. Invest. Drugs* **2000**, *9*, 2393-2402.
9. Packer, L.; Witt, E. H.; Tritschler, H. J. alpha-Lipoic acid as a biological antioxidant. *Free Radical Biol. Med.* **1995**, *19*, 227-250.
10. Tirosh, O.; Sen, C. K.; Roy, S.; Kobayashi, M. S.; Packer, L. Neuroprotective effects of alpha-lipoic acid and its positively charged amide analogue. *Free Radical Biol. Med.* **1999**, *26*, 1418-1426.
11. Biewenga, G. P.; Haenen, G. R.; Bast, A. The pharmacology of the antioxidant lipoic acid. *Gen. Pharmacol.* **1997**, *29*, 315-331.
12. Packer, L. alpha-Lipoic acid: a metabolic antioxidant which regulates NF-kappaB signal transduction and protects against oxidative injury. *Drug Metab. Rev.* **1998**, *30*, 245-275.
13. Zhang, L.; Xing, G. Q.; Barker, J. L.; Chang, Y.; Maric, D.; Ma, W.; Li, B. S.; Rubinow, D. R. alpha-Lipoic acid protects rat cortical neurons against cell death induced by amyloid and hydrogen peroxide through the Akt signalling pathway. *Neurosci. Lett.* **2001**, *312*, 125-128.
14. Hager, K.; Marahrens, A.; Kenklies, M.; Riederer, P.; Munch, G. alpha-Lipoic acid as a new treatment option for Alzheimer type dementia. *Arch. Gerontol. Geriatr.* **2001**, *32*, 275-282.

15. Inestrosa, N. C.; Alvarez, A.; Perez, C. A.; Moreno, R. D.; Vicente, M.; Linker, C.; Casanueva, O. I.; Soto, C.; Garrido, J. Acetylcholinesterase accelerates assembly of amyloid-beta-peptides into Alzheimer's fibrils: possible role of the peripheral site of the enzyme. *Neuron* **1996**, *16*, 881-891.
16. Bartolini, M.; Bertucci, C.; Cavrini, V.; Andrisano, V.  $\beta$ -Amyloid aggregation induced by human acetylcholinesterase: inhibition studies. *Biochem. Pharmacol.* **2003**, *65*, 407-416.
17. Ellman, G. L.; Courtney, K. D.; Andres, V.; Featherstone, R. M. A new rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* **1961**, *7*, 88-95.
18. Recanatini, M.; Cavalli, A.; Belluti, F.; Piazzzi, L.; Rampa, A.; Bisi, A.; Gobbi, S.; Valenti, P.; Andrisano, V.; Bartolini, M.; Cavrini, V. SAR of 9-amino-1,2,3,4-tetrahydroacridine-based acetylcholinesterase inhibitors: synthesis, enzyme inhibitory activity, QSAR, and structure-based CoMFA of tacrine analogues. *J. Med. Chem.* **2000**, *43*, 2007-2018.
19. Pang, Y. P.; Quiram, P.; Jelacic, T.; Hong, F.; Brimijoin, S. Highly potent, selective, and low cost bis-tetrahydroaminacrine inhibitors of acetylcholinesterase. Steps toward novel drugs for treating Alzheimer's disease. *J. Biol. Chem.* **1996**, *271*, 23646-23649.
20. Inestrosa, N. C.; Alvarez, A.; Calderon, F. Acetylcholinesterase is a senile plaque component that promotes assembly of amyloid  $\beta$ -peptide into Alzheimer's filaments. *Mol. Psychiatry* **1996**, *1*, 359-361.
21. Piazzzi, L.; Rampa, A.; Bisi, A.; Gobbi, S.; Belluti, F.; Cavalli, A.; Bartolini, M.; Andrisano, V.; Valenti, P.; Recanatini, M. 3-(4-[[Benzyl(methyl)amino]methyl]phenyl)-6,7-dimethoxy-2H-2-chromenone (AP2238) inhibits both acetylcholinesterase and acetylcholinesterase-induced  $\beta$ -amyloid aggregation: a dual function lead for Alzheimer's disease therapy. *J. Med. Chem.* **2003**, *46*, 2279-2282.
22. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **1983**, *65*, 55-63.

Table 1. Inhibition of AChE and BChE Activities by 15, Tacrine (TC), Lipoic Acid (LA), and Hybrid Derivatives 1-8 (See Figure 1 for Structures)

| compd | n | R  | IC <sub>50</sub> ± SEM (nM) <sup>a</sup> |             |
|-------|---|----|------------------------------------------|-------------|
|       |   |    | AChE                                     | BChE        |
| 1     | 2 | H  | 97.0 ± 3.6                               | 47.5 ± 1.8  |
| 2     | 3 | H  | 6.96 ± 0.45                              | 12.0 ± 0.6  |
| 3     | 4 | H  | 35.2 ± 2.2                               | 5.04 ± 0.32 |
| 4     | 5 | H  | 38.4 ± 2.3                               | 1.48 ± 0.35 |
| 5     | 6 | H  | 30.1 ± 1.5                               | 3.24 ± 0.29 |
| 6     | 7 | H  | 32.7 ± 1.3                               | 8.58 ± 0.57 |
| 7     | 3 | Cl | 0.253 ± 0.016                            | 10.8 ± 2.5  |
| 8     |   |    | 1090 ± 180                               | 329 ± 28    |
| 15    |   |    | 21.5 ± 0.8                               | 2580 ± 60   |
| TC    |   |    | 424 ± 21                                 | 45.8 ± 3.0  |
| LA    |   |    | >1000000                                 | >1000000    |

<sup>a</sup> Human recombinant AChE and BChE from human serum were used. IC<sub>50</sub> values represent the concentration of inhibitor required to decrease enzyme activity by 50% and are the mean of two independent measurements, each performed in triplicate. See Supporting Information for details.

Table 2. Effects of the Hybrid Compound 7 and the Two Parent Compounds LA and 15 on Intracellular ROS Formation in Neuronal Cells<sup>a</sup>

| concn, $\mu$ M | intracellular ROS, % |                |                  |
|----------------|----------------------|----------------|------------------|
|                | LA                   | 7              | 15               |
| 0              | 86.00 ± 9.46         | 86.00 ± 9.46   | 86.00 ± 9.46     |
| 0.1            | 91.25 ± 2.99         | 88.75 ± 8.41   | 99.50 ± 3.54     |
| 0.5            | 90.25 ± 6.65         | 95.00 ± 5.20   | 99.67 ± 7.02     |
| 1              | 83.00 ± 7.58         | 77.00 ± 8.76   | 96.67 ± 8.02     |
| 5              | 79.50 ± 8.06         | 61.67 ± 9.02*  | 82.50 ± 7.78     |
| 10             | 74.33 ± 3.51         | 51.25 ± 9.21** | tox <sup>b</sup> |

|    |               |                |                  |
|----|---------------|----------------|------------------|
| 50 | 58.50 ± 9.19* | 30.50 ± 9.04** | tox <sup>b</sup> |
|----|---------------|----------------|------------------|

<sup>a</sup> The results are expressed as percentage increase of intracellular ROS evoked by exposure to *tert*-butyl hydroperoxide. Values are the mean ± SD of three independent experiments (treated vs untreated; (\*)  $p < 0.01$ , (\*\*)  $p < 0.001$ ).<sup>b</sup> tox = cytotoxicity. See Supporting Information for details.

### 1. Synthesis and characterization of compounds 1-8

Compounds 1-8 were synthesized according to Scheme 1, coupling tetrahydroacridine intermediates with lipoic acid.

Melting points were taken in glass capillary tubes on a Büchi SMP-20 apparatus and are uncorrected. IR, electron impact (EI) mass, and direct infusion ESI-MS spectra were recorded on Perkin-Elmer 297, VG 7070E, and Waters ZQ 4000 apparatus, respectively. <sup>1</sup>H NMR, <sup>13</sup>C NMR, gHSQC and COSY experiments were recorded on Mercury 400 and Varian VXR 200 and 300 instruments. Chemical shifts are reported in parts per million (ppm) relative to tetramethylsilane (TMS), and spin multiplicities are given as s (singlet), d (doublet), dd (double doublet), t (triplet), or m (multiplet). Although the IR spectra data are not included (because of the lack of unusual features), they were obtained for all compounds reported and were consistent with the assigned structures. The elemental compositions of the compounds agreed to within ± 0.4% of the calculated value. When the elemental analysis is not included, crude compounds were used in the next step without further purification. Chromatographic separations were performed on silica gel columns by flash (Kieselgel 40, 0.040-0.063 mm; Merck) or gravity column (Kieselgel 60, 0.063-0.200 mm; Merck) chromatography. Compounds were named following IUPAC rules as applied by Beilstein-Institut AutoNom (version 2.1), a PC integrated software package for systematic names in organic chemistry.

**(3-Aminomethyl-6-chloro-1,2,3,4-tetrahydroacridin-9-yl)amine (16).** The synthesis of compound 16 was achieved by condensation of 2-amino-4-chlorobenzonitrile with 3-nitromethylcyclohexanone followed by reduction of the nitro group according to Rosini et al.,<sup>1</sup> and the isomeric conformation was assigned by means of <sup>1</sup>H NMR, <sup>13</sup>C NMR, gHSQC, and COSY experiments. Total yield 30%; mp 285-288 °C; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.91 (d,  $J = 8.9$  Hz, 1H, C8-H), 7.58 (d,  $J = 2.3$  Hz, 1H, C5-H), 7.19 (dd,  $J = 9.0, 2.3$  Hz, 1H, C7-H), 2.86-2.94 (m, 1H, C4-H), 2.60-2.69 (m, 3H, -CH<sub>2</sub>NH<sub>2</sub>, C1-H), 2.19-2.25 (m, 2H, C1-H, C4-H), 2.04-2.13 (m, 1H, C2-H), 1.75-1.83 (m, 1H, C3-H), 1.29-1.39 (m, 1H, C2-H); <sup>13</sup>C NMR (100 MHz, CD<sub>3</sub>OD) δ 158.9, 150.3, 147.7, 135.2, 126.1 (C5), 124.6 (C7), 124.1 (C8), 116.3, 110.4, 48.1 (-CH<sub>2</sub>NH<sub>2</sub>), 38.3 (C4), 37.9 (C3), 27.3 (C2), 24.2 (C1); EI MS  $m/z$  261 (M<sup>+</sup>).



### General procedure for the synthesis of compounds 1-8.

A solution of the appropriate tetrahydroacridinamine (1 eq) and lipoic acid (1.5 eq) in dry DMF (5 mL), under N<sub>2</sub>, was cooled to 0 °C and then treated with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI) (1.2 eq): the mixture was stirred at 0 °C for further 15 min and then at rt for 2 h in the dark. Solvent was then removed under vacuum, avoiding heating up the reaction mixture, affording an oily residue that was purified by gravity column.

#### 5-([1,2]Dithiolan-3-yl)-*N*-{[2-(1,2,3,4-tetrahydroacridin-9-yl)amino]ethyl}pentanamide (1).

It was synthesized from *N*1-(1,2,3,4-tetrahydroacridin-9-yl)ethane-1,2-diamine (9)<sub>2</sub> (140 mg).

Elution with petroleum ether/CH<sub>2</sub>Cl<sub>2</sub>/MeOH/aqueous 30% ammonia (6:3:1:0.055) afforded 1 as a foam solid: 35% yield; <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.12 (d, *J* = 8.8 Hz, 1H), 7.78 (d, *J* = 8.8 Hz, 1H), 7.58, (t, *J* = 8.2 Hz, 1H), 7.39 (t, *J* = 8.2 Hz, 1H), 3.70 (t, *J* = 6.3 Hz, 2H), 3.28-3.39 (m, 3H), 2.93-3.15 (m, 4H), 2.71-2.79 (m, 2H), 2.26-2.40 (m, 1H), 2.15 (t, *J* = 8.6 Hz, 2H), 1.64-1.93 (m, 5H), 1.30-1.61 (m, 6H); MS (ESI<sup>+</sup>) *m/z* 430 (M+H)<sup>+</sup>. Calcd. for C<sub>23</sub>H<sub>31</sub>N<sub>3</sub>OS<sub>2</sub>: C, 64.30; H, 7.27; N, 9.78; found C, 64.41; H, 7.28; N, 9.75.

#### 5-([1,2]Dithiolan-3-yl)-*N*-{[3-(1,2,3,4-tetrahydroacridin-9-yl)amino]propyl}pentanamide (2).

It was synthesized from *N*1-(1,2,3,4-tetrahydroacridin-9-yl)propane-1,3-diamine (10) (100 mg, obtained from 9-chloro-1,2,3,4-tetrahydroacridine and propane-1,3-diamine following the procedure described in Carlier et al.<sup>3</sup> and purified by flash chromatography with a step gradient system of CH<sub>2</sub>Cl<sub>2</sub>/MeOH/ aqueous 30% ammonia (9.5:0.5:0.0 to 7:3:0.1): 65% yield, <sup>1</sup>H NMR (200 MHz, CD<sub>3</sub>OD) δ 8.08 (d, *J* = 8.8 Hz, 1H), 7.78 (d, *J* = 8.7 Hz, 1H), 7.53, (t, *J* = 8.3 Hz, 1H), 7.32 (t, *J* = 8.3 Hz, 1H), 3.54 (t, *J* = 6.7 Hz, 2H), 2.87-2.98 (m, 2H), 2.65 (t, *J* = 7.5 Hz, 4H), 1.64-1.93 (m, 6H). Elution with petroleum ether/CH<sub>2</sub>Cl<sub>2</sub>/MeOH/aqueous 30% ammonia (5:4:1:0.05) afforded 2 as a foam solid: 35% yield; <sup>1</sup>H NMR (200 MHz, CD<sub>3</sub>OD) δ 8.15 (d, *J* = 8.8 Hz, 1H), 7.78 (d, *J* = 8.8 Hz, 1H), 7.56-7.64 (m, 1H), 7.37-7.44 (m, 1H), 3.69 (t, *J* = 6.6 Hz, 2H), 3.40-3.52 (m, 1H), 3.23-3.36 (t, *J* = 6.6 Hz, 2H), 2.92-3.18 (m, 4H), 2.74-2.83 (m, 2H), 2.28-2.43 (m, 1H), 2.19 (t, *J* = 7.1 Hz, 2H), 1.73-1.95 (m, 7H), 1.22-1.68 (m, 6H); MS (ESI<sup>+</sup>) *m/z* 444 (M+H)<sup>+</sup>. Calcd. for C<sub>24</sub>H<sub>33</sub>N<sub>3</sub>OS<sub>2</sub>: C, 64.97; H, 7.50; N, 9.47; found C, 65.18; H, 7.52; N, 9.44.

#### 5-([1,2]Dithiolan-3-yl)-*N*-{[4-(1,2,3,4-tetrahydroacridin-9-yl)amino]butyl}pentanamide (3).

It was synthesized from *N*1-(1,2,3,4-tetrahydroacridin-9-yl)butane-1,4-diamine (11)<sub>3</sub> (290 mg). Elution with petroleum ether/CH<sub>2</sub>Cl<sub>2</sub>/MeOH/aqueous 30% ammonia (6:3:1:0.06) afforded 3 as a foam solid: 38% yield; <sup>1</sup>H NMR (200 MHz, CD<sub>3</sub>OD) δ 8.12 (d, *J* = 8.6 Hz, 1H), 7.78 (d, *J* = 8.6 Hz, 1H), 7.52-7.62 (m, 1H), 7.32-7.43 (m, 1H), 3.41-3.60 (m, 3H), 2.90-3.21 (m, 6H), 2.68-2.77 (m, 2H), 2.31-2.46 (m, 1H), 2.17 (t, *J* = 6.9 Hz, 2H), 1.38-1.95 (m, 15H); MS (ESI<sup>+</sup>) *m/z* 458 (M+H)<sup>+</sup>.

Calcd. for  $C_{25}H_{35}N_3OS_2$ : C, 65.60; H, 7.71; N, 9.18; found C, 65.67; H, 7.69; N, 9.15.

**5-([1,2]Dithiolan-3-yl)-N-{[5-(1,2,3,4-tetrahydroacridin-9-yl)amino]pentyl}pentanamide**

(4). It was synthesized from *N*1-(1,2,3,4-tetrahydroacridin-9-yl)pentane-1,5-diamine (12)<sub>3</sub> (480 mg). Elution with petroleum ether/ $CH_2Cl_2$ /MeOH/aqueous 30% ammonia (6:3:1:0.055) afforded 4 as a foam solid: 40% yield;  $^1H$  NMR (200 MHz,  $CD_3OD$ )  $\delta$  8.09 (d,  $J$  = 8.6 Hz, 1H), 7.78 (d,  $J$  = 8.6 Hz, 1H), 7.52-7.60 (m, 1H), 7.33-7.41 (m, 1H), 3.40-3.57 (m, 3H), 2.87-3.18 (m, 6H), 2.63-2.75 (m, 2H), 2.25-2.43 (m, 1H), 2.17 (t,  $J$  = 6.8 Hz, 2H), 1.35-1.95 (m, 17H); MS (ESI<sup>+</sup>)  $m/z$  472 (M+H)<sup>+</sup>. Calcd. for  $C_{26}H_{37}N_3OS_2$ : C, 66.20; H, 7.91; N, 8.91; found C, 66.41; H, 7.89; N, 8.88.

**5-([1,2]Dithiolan-3-yl)-N-{[6-(1,2,3,4-tetrahydroacridin-9-yl)amino]hexyl}pentanamide (5).**

It was synthesized from *N*1-(1,2,3,4-tetrahydroacridin-9-yl)hexane-1,6-diamine (13)<sub>3</sub> (370 mg). Elution with petroleum ether/ $CH_2Cl_2$ /MeOH/aqueous 30% ammonia (6:3:1:0.05) afforded 5 as a foam solid: 30% yield;  $^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.83 (apparent t,  $J$  = 9.3 Hz, 2H), 7.47-7.56 (m, 1H), 7.28-7.37 (m, 1H), 5.89 (t,  $J$  = 3.2 Hz, 1H, exchangeable with  $D_2O$ ), 4.15 (br s, 2H, exchangeable with  $D_2O$ ), 3.40-3.57 (m, 3H), 3.01-3.23 (m, 6H), 2.60-2.75 (m, 2H), 2.31-2.48 (m, 1H), 2.15 (t,  $J$  = 7.3 Hz, 2H), 1.35-1.96 (m, 19H); MS (ESI<sup>+</sup>)  $m/z$  486 (M+H)<sup>+</sup>. Calcd. for  $C_{27}H_{39}N_3OS_2$ : C, 66.76; H, 8.09; N, 8.65; found C, 66.87; H, 8.12; N, 8.62.

S5

**5-([1,2]Dithiolan-3-yl)-N-{[7-(1,2,3,4-tetrahydroacridin-9-yl)amino]heptyl}pentanamide (6).**

It was synthesized from *N*1-(1,2,3,4-tetrahydroacridin-9-yl)heptane-1,7-diamine (14)<sub>3</sub> (220 mg). Elution with petroleum ether/ $CH_2Cl_2$ /MeOH/aqueous 30% ammonia (6:3:1:0.05) afforded 6 as a foam solid: 35% yield;  $^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.92 (apparent t,  $J$  = 9.4 Hz, 2H), 7.51-7.61 (m, 1H), 7.30-7.41 (m, 1H), 5.57 (t,  $J$  = 3.2 Hz, 1H, exchangeable with  $D_2O$ ), 3.40-3.61 (m, 3H), 3.01-3.24 (m, 6H), 2.64-2.73 (m, 2H), 2.38-2.54 (m, 1H), 2.18 (t,  $J$  = 7.3 Hz, 2H), 1.25-1.98 (m, 21H); MS (ESI<sup>+</sup>)  $m/z$  500 (M+H)<sup>+</sup>. Calcd. for  $C_{28}H_{41}N_3OS_2$ : C, 67.29; H, 8.27; N, 8.41; found C, 67.43; H, 8.30; N, 8.39.

**5-([1,2]Dithiolan-3-yl)-N-[3-(6-chloro-1,2,3,4-tetrahydro-acridin-9-yl)amino]propyl}pentanamide (7).**

It was synthesized from *N*1-(6-chloro-1,2,3,4-tetrahydroacridin-9-yl)propane-1,3-diamine (15) (180 mg, obtained from 6,9-dichloro-1,2,3,4-tetrahydroacridine and propane-1,3-diamine following the procedure described in Carlier et al.,<sup>3</sup> and purified by flash chromatography with a step gradient system of  $CH_2Cl_2$ /MeOH/ aqueous 30% ammonia (9.5:0.5:0.0 to 8:2:0.03): 70% yield,  $^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.93 (d,  $J$  = 9.1 Hz, 1H), 7.86 (d,  $J$  = 2.4 Hz, 1H), 7.22 (dd,  $J$  = 9.0, 2.3 Hz, 1H), 3.62 (t,  $J$  = 6.8 Hz, 2H), 2.88-3.05 (m, 4H), 2.60-2.68 (m, 2H), 1.71-1.95 (m, 6H). Elution with petroleum ether/ $CH_2Cl_2$ /EtOH/aqueous 30% ammonia (7:2:1:0.03) afforded 7 as a foam solid: 35% yield;  $^1H$  NMR (200 MHz,  $CD_3OD$ )  $\delta$

8.08 (d,  $J = 8.9$  Hz, 1H), 7.72 (d,  $J = 2.1$  Hz, 1H), 7.28 (dd,  $J = 8.9, 2.1$  Hz, 1H), 3.42-3.58 (m, 3H), 3.27 (t,  $J = 6.5$  Hz, 2H), 2.89-3.17 (m, 4H), 2.65-2.77 (m, 2H), 2.27-2.43 (m, 1H), 2.19 (t,  $J = 7.2$  Hz, 2H), 1.73-1.91 (m, 7H), 1.31-1.65 (m, 6H); MS (ESI+)  $m/z$  478 (M+H)<sup>+</sup>. Calcd. for C<sub>24</sub>H<sub>32</sub>ClN<sub>3</sub>OS<sub>2</sub>: C, 60.29; H, 6.75; N, 8.79; found C, 60.45; H, 6.74; N, 8.77.

***N*-[*(9-Amino-6-chloro-1,2,3,4-tetrahydroacridin-3-yl)methyl*]-5-[1,2]dithiolan-3-yl)pentanamide (8).** It was synthesized from **1** **6** (150 mg). Elution with

CH<sub>2</sub>Cl<sub>2</sub>/toluene/EtOH/aqueous 30% ammonia (5:3:2:0.02) afforded **8** as a foam solid: 30% yield; <sup>1</sup>H NMR (200 MHz, CD<sub>3</sub>OD)  $\delta$  8.09 (d,  $J = 8.9$  Hz, 1H), 7.72 (d,  $J = 2.2$  Hz, 1H), 7.36 (dd,  $J = 9.2, 2.2$  Hz, 1H), 3.50-3.62 (m, 2H), 2.96-3.21 (m, 4H), 2.70-2.83 (m, 1H), 2.38-2.69 (m, 3H), 2.28 (t,  $J = 7.0$  Hz, 2H), 2.05-2.21 (m, 2H), 1.79-1.95 (m, 1H), 1.23-1.78 (m, 7H); EI MS  $m/z$  449 (M<sup>+</sup>). Calcd. for C<sub>22</sub>H<sub>28</sub>ClN<sub>3</sub>OS<sub>2</sub>: C, 58.71; H, 6.27; N, 9.34; found C, 58.91; H, 6.26; N, 9.31.

## 2. Biology

**Inhibition of AChE and BChE.** The method of Ellman et al. was followed.<sup>4</sup> Five different concentrations of each compound were used in order to obtain inhibition of AChE or BChE activity comprised between 20-80%. The assay solution consisted of a 0.1 M phosphate buffer pH 8.0, with the addition of 340  $\mu$ M 5,5'-dithio-bis(2-nitrobenzoic acid), 0.02 unit/mL of human recombinant AChE or human serum BChE (Sigma Chemical), and 550  $\mu$ M of substrate (acetylthiocholine iodide or butyrylthiocholine iodide). Test compounds were added to the assay solution and preincubated at 37 °C with the enzyme for 20 min followed by the addition of substrate. Assays were done with a blank containing all components except AChE or BChE in order to account for non-enzymatic reaction. The reaction rates were compared and the percent inhibition due to the presence of test compounds was calculated. Each concentration was analyzed in triplicate, and IC<sub>50</sub> values were determined graphically from log concentration-inhibition curves.

**Determination of Steady State Inhibition Constant.** To obtain estimates of the competitive inhibition constant  $K_i$ , reciprocal plots of  $1/V$  versus  $1/[S]$  were constructed at relatively low concentration of substrate (below 0.5 mM). The plots were assessed by a weighted least square analysis that assumed the variance of  $V$  to be a constant percentage of  $V$  for the entire data set. Slopes of these reciprocal plots were then plotted against the concentration of **7** (range 0 – 0.344 nM) in a weighted analysis and  $K_i$  was determined as the ratio of the replot intercept to the replot slope.

Reciprocal plots involving **TC** (not shown) or **7** inhibition show both increasing slopes (decreased  $V_{max}$  at increasing inhibitor's concentrations) and increasing intercepts (higher  $K_m$ ) with higher inhibitor concentration. This pattern indicates mixed inhibition, arising from significant

inhibitor interaction with both the free enzyme and the acetylated enzyme. Replots of the slope  
S7

versus the concentration of 7 or TC gives estimate of competitive inhibition constant,  $K_i = 0.155 \pm 0.046$  nM or  $K_i = 0.151 \pm 0.016$   $\mu$ M, respectively.

So the pattern in the graphical representation shows 7 able to bind to the peripheral anionic site as well as the active site of AChE.

**Inhibition of AChE-induced A  $\beta$  aggregation.** Aliquots of 2  $\mu$ L A  $\beta$  peptide, lyophilized from 2 mg mL<sup>-1</sup> 1,1,1,3,3,3-hexafluoro-2-propanol solution and dissolved in DMSO, were incubated for 24 h at room temperature in 0.215 M sodium phosphate buffer (pH 8.0) at a final concentration of 230  $\mu$ M. For co-incubation experiments aliquots (16  $\mu$ L) of AChE (final concentration 2.30  $\mu$ M, A $\beta$ /AChE molar ratio 100:1) and AChE in the presence of 2  $\mu$ L of the tested inhibitor in 0.215 M sodium phosphate buffer pH 8.0 solution (final inhibitor concentration 100  $\mu$ M) were added. Blanks containing A  $\beta$ , AChE, and A  $\beta$  plus inhibitors at various concentrations, in 0.215 M sodium phosphate buffer (pH 8.0) were prepared. The final volume of each vial was 20  $\mu$ L. Each assay was run in duplicate. To quantify amyloid fibril formation, the thioflavin T (ThT) fluorescence method was then applied.<sup>5</sup> After dilution with glycine-NaOH buffer (pH 8.5), containing 1.5 mM ThT, the fluorescence intensities due to  $\beta$ -sheet conformation was monitored for 300 s at  $\lambda_{em} = 490$  nm ( $\lambda_{ex} = 446$  nm). The percent inhibition of the AChE induced aggregation due to the presence of the test compound was calculated by the following expression:  $100 - (IF_i/IF_o \times 100)$  where  $IF_i$  and  $IF_o$  are the fluorescence intensities obtained for A $\beta$  plus AChE in the presence and in the absence of inhibitor, respectively, minus the fluorescent intensities due to the respective blanks.

**Cell cultures.** Human neuronal-like cells, SH-SY5Y, were routinely grown at 37°C in a humidified incubator with 5% CO<sub>2</sub> in Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum (FCS), 2 mM glutamine, 50 U/ml penicillin and 50  $\mu$ g/ml streptomycin.

**Determination of cytotoxicity.** The cytotoxicity was evaluated with the colorimetric MTT [3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide] assay, as described by Mosmann et al.<sup>6</sup> Briefly, SH-SY5Y cells were seeded in 96-well microtiter plates at  $2 \times 10^5$  cells/well. After  
S8

24 h of incubation at 37°C in 5% CO<sub>2</sub>, the growth medium was removed and media containing compounds (0.1-50  $\mu$ M) were added to the cells. After 24 h of incubation, the cells were washed with phosphate buffered saline (PBS) and then incubated with MTT (5 mg/ml) in PBS for 4 h. After removal of MTT and further washing, the formazan crystals were dissolved with isopropanol. The amount of formazan was measured (405 nm) with a spectrophotometer (TECAN<sup>®</sup>, Spectra model Classic, Salzburg, Austria). The cell viability was expressed as percentage of control cells and

calculated by the formula  $F_t/F_{nt} \times 100$ , where  $F_t$  = absorbance of treated neurones and  $F_{nt}$  = absorbance of untreated neurones.

**Determination of antioxidant activity.** The antioxidant activity of compounds was evaluated by measuring the formation of intracellular reactive oxygen species (ROS) evoked by exposure of SH-SY5Y cells to *tert*-butyl hydroperoxide (*t*-BuOOH), a compound used to induce oxidative stress. Formation of intracellular ROS was determined using a fluorescent probe, DCFH-DA, as described by Wang H. et al.<sup>7</sup> Briefly, SH-SY5Y cells were seeded in 96-well microtiter plates at  $2 \times 10^5$  cells/well. After 24 h of incubation at 37°C in 5% CO<sub>2</sub>, the growth medium was removed and media containing compounds (0.1-50 µM) were added to the cells. After 24 h of incubation, the cells were washed with PBS and then incubated with 5 µM of DCFH-DA in PBS at 37°C in 5% CO<sub>2</sub> for 30 min. After removal of DCFH-DA and further washing, the cells were incubated with 0.1 mM *t*-BuOOH in PBS for 30 min. At the end of incubation, the fluorescence of the cells from each well was measured ( $\lambda_{\text{excitation}} = 485 \text{ nm}$ ,  $\lambda_{\text{emission}} = 535 \text{ nm}$ ) with a spectrofluorometer (Wallac Victor Multilabel Counter, Perkin Elmer Inc., Boston, MA). The results were expressed as percentage increase of intracellular ROS evoked by exposure to *t*-BuOOH and calculated by the formula  $[(F_t - F_{nt}) / F_{nt} \times 100]$ , where  $F_t$  = fluorescence of treated neurones and  $F_{nt}$  = fluorescence of untreated neurones.

**Statistical analysis.** Data are reported as mean  $\pm$  SD of at least 3 independent experiments.

Statistical analysis was performed using ANOVA (Scheffe post hoc test was used) and the S9


differences were considered significant at  $p < 0.05$ . Analyses were performed using STATISTICA 4.5 software on a Windows platform.

## References

- (1) Rosini, M.; Antonello, A.; Cavalli, A.; Bolognesi, M. L.; Minarini, A.; Marucci, G.; Poggesi, E.; Leonardi, A.; Melchiorre, C. Prazosin-related compounds. Effect of transforming the piperazinylquinazoline moiety into an aminomethyltetrahydroacridine system on the affinity for  $\alpha_1$ -adrenoreceptors. *J. Med. Chem.* **2003**, *46*, 4895-4903.
- (2) Steinberg, G. M.; Mednick, M. L.; Maddox, J.; Rice, R. A hydrophobic binding site in acetylcholinesterase. *J. Med. Chem.* **1975**, *18*, 1057-1061.
- (3) Carlier, P. R.; Du, D. M.; Han, Y.; Liu, J.; Pang, Y. P. Potent, easily synthesized huperzine A-tacrine hybrid acetylcholinesterase inhibitors. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 2335-2338.
- (4) Ellman, G. L.; Courtney, K. D.; Andres, V.; Featherstone, R. M. A new rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* **1961**, *7*, 88-95.


- (5) Bartolini, M.; Bertucci, C.; Cavrini, V.; Andrisano, V.  $\beta$ -Amyloid aggregation induced by human acetylcholinesterase: inhibition studies. *Biochem. Pharmacol.* **2003**, *65*, 407-416.
- (6) Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **1983**, *65*, 55-63.
- (7) Wang, H.; Joseph, J. A. Quantifying cellular oxidative stress by dichlorofluorescein assay using microplate reader. *Free Radic. Biol. Med.* **1999**, *27*, 612-616.

The general structure of lipoic acid derivatives can be schematized as such:



Where  $n = 1-4$ ;

Examples where R =:




The examples are structures (1-8) reported in the enclosed manuscript;

An other example is an inferior homologous of 8:

**RM11**:  $n = 1$ ,  $\text{R}_1 = 3\text{-ylmethylamine}$ ,  $\text{R}_2 = \text{NH}_2$ ,  $\text{R}_3 = 6\text{-chloro}$ ;


b)



Example:

**MR106:**  $n = 4$ ;  $X = \text{CH=CH}$ ,  $R_1 = 2\text{-MeO}$ ,  $m = 6$ ;  $R_2 = \text{Et}$ ,  $R_3 = \text{H}$ .

c)



$R_1 = \text{alkyl}$ ,  $R_2 = \text{alkyl}$ , alkylamine

$Y = \text{alkyl}$

$X = \text{OH}$ , O-carbamoyl, O-


An example is **MR112**:  $Y = \text{MeCH}$ ,  $R_1 = R_2 = \text{Me}$ ,  $X = \text{O-}$ .

Synthesis of examples:

**N-(9-Amino-6-chloro-1,2,3,4-tetrahydro-acridin-3-yl-methyl)-2-[1,2]dithiolan-3-yl-acetamide (RM11).** A solution of 3-aminomethyl-6-chloro-1,2,3,4-tetrahydro-acridin-9-ylamine<sup>i</sup> (140 mg, 0.53 mmol) and [1,2]Dithiolan-3-yl-acetic acid<sup>ii</sup> (90 mg, 0.55 mmol) in dry DMF (5 mL), under  $\text{N}_2$ , was cooled to  $0^\circ\text{C}$  and then added of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI) (113 mg, 0.59 mmol). The mixture was stirred at  $0^\circ\text{C}$  for further 15 min and then at rt for 2 h in the dark. Solvent was then removed under vacuum, avoiding heating up the reaction mixture, affording a residue that was purified by gravity column. Elution with petroleum ether/ $\text{CH}_2\text{Cl}_2$ /MeOH/aqueous 30% ammonia (5:4:1:0.1) afforded **RM11** as a foam solid: 40% yield;  $^1\text{H}$  NMR (200 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.03 (d,  $J = 9.2$  Hz, 1H), 7.67 (d,  $J = 2.2$  Hz, 1H), 7.30 (dd,  $J = 8.8$ , 2.2 Hz, 1H), 4.05 (m, 1H), 2.95-3.34 (m, 5H), 2.46-2.59 (m, 6H), 1.99-2.20 (m, 3H), 1.48-1.52 (m, 1H). Anal.  $\text{C}_{19}\text{H}_{22}\text{ClN}_3\text{OS}_2$ : C, 55.93; H, 5.44; N, 10.30. Found: C, 56.01; H, 5.45, N, 10.11.

**5-[1,2]Dithiolan-3-yl-pentanoic acid {6-[Ethyl-(2-methoxy-benzyl)-amino]-esyl}-amide (MR106)** was synthesized from  $N^1$ -ethyl- $N^1$ -(2-methoxy-benzyl)-hexano-1,6-diamine<sup>iii</sup> (300 mg, 1.13 mmol) and lipoic acid (350 mg, 1.70 mmol) according to the procedure described for **RM11**, and purified by gravity column. Elution with a gradient of mobile phase petroleum ether/toluene/ $\text{CH}_2\text{Cl}_2$ /EtOH/aqueous 30% ammonia (from 7:2:1:1:0.05 to 7:1:1:1:0.05) afforded **MR106** as solid foam: yield 43%,  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.42-7.48 (m, 1H); 7.18-7.26 (m, 1H), 6.85-6.99 (m, 2H), 5.43 (br t, 1H, exchangeable with  $\text{D}_2\text{O}$ ), 3.84 (s, 3H), 3.53-3.64 (m, 1H + s, 2H), 3.08-3.27 (m, 4H), 2.43-2.57 (m, 5H), 2.17 (t,  $J = 7.4$  Hz, 2H), 1.82-2.00 (m, 1H), 1.29-1.73 (m, 14H), 1.07 (t,  $J = 7.0$  Hz, 3H); MS (ESI<sup>+</sup>)  $m/z$  453 (M+H)<sup>+</sup>. Calculated Anal. for  $\text{C}_{24}\text{H}_{40}\text{N}_2\text{O}_2\text{S}_2$ : C, 63.67; H, 8.91; N, 6.19. Found: C, 63.79; H, 8.93, N, 6.17.

**{2-[3-(1-dimethylamino-ethyl)-phenoxy]-ethyl}-carbamic acid Tert-butyl ester (II).** A solution of **I<sup>iv</sup>** (350 mg, 2.17 mmol), (3-chloro-propyl)-carbamic acid tert-butylic ester (420 mg, 2.17 mmol) and  $K_2CO_3$  (300 mg, 2.17 mmol) in DMF (10 ml) was stirred under reflux for 24 h. Solvent evaporation afforded a residue that was purified by gravity column. Elution with  $CHCl_3/MeOH/$  aqueous 30% ammonia (9:1:0.02) afforded **II** as an oil: yield 65%,  $^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.20 (t,  $J = 8.0$  Hz, 1H); 6.79-6.89 (m, 3H), 4.92 (br s, 1H, exchangeable with  $D_2O$ ), 4.02 (t,  $J = 6.4$  Hz, 2H); 3.20-3.33 (m, 3H), 2.20 (s, 6H), 1.93-1.99 (m, 2H), 1.44 (s, 9H), 1.35 (d,  $J = 6.6$  Hz, 3H).



**{2-[3-(1-dimethylamino-ethyl)-phenoxy]-ethyl}-carbamic acid tert-butyl ester (II).** A solution of **I<sup>iv</sup>** (350 mg, 2.17 mmol), (3-chloro-propyl)-carbamic acid tert-butylic ester (420 mg, 2.17 mmol) and  $K_2CO_3$  (300 mg, 2.17 mmol) in DMF (10 ml) was stirred under reflux for 24 h. Solvent evaporation afforded a residue that was purified by gravity column. Elution with  $CHCl_3/MeOH/$  aqueous 30% ammonia (9:1:0.02) afforded **II** as an oil: yield 65%,  $^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.20 (t,  $J = 8.0$  Hz, 1H); 6.79-6.89 (m, 3H), 4.92 (br s, 1H, exchangeable with  $D_2O$ ), 4.02 (t,  $J = 6.4$  Hz, 2H); 3.20-3.33 (m, 3H), 2.20 (s, 6H), 1.93-1.99 (m, 2H), 1.44 (s, 9H), 1.35 (d,  $J = 6.6$  Hz, 3H).

**3-[3-(1-Dimethylamino-ethyl)-phenoxy]-propylamine (III).** A solution of **II** (200 mg, 0.62 mmol) in  $CH_2Cl_2$  (5 mL) is added with TFA (1.5 mL) and stirred at room temperature for 2 h. The reaction mixture was evaporated under vacuum, the afforded residue was dissolved in water, resobasico by adding NaOH 2 N and then extracted with  $CHCl_3$  (3 x 20 mL). The evaporation of the dried solvent afforded **III** as an oil; quantitative yield,  $^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.20 (t,  $J = 8.0$



Hz, 1H); 6.72-6.88 (m, 3H), 4.04 (t,  $J = 6.2$  Hz, 2H); 3.12-3.22 (m, 1H), 2.91 (t,  $J = 6.6$  Hz, 2H), 2.19 (s, 6H), 1.88-1.95 (m, 2H), 1.43 (br s, 2H, exchangeable with  $D_2O$ ), 1.34 (d,  $J = 6.6$  Hz, 3H).

**5-[1,2]Dithiolan-3-yl-pentanoic acid {3-[3-(1-Dimethylamino-ethyl)-phenoxy]-propyl}-amide (MR112)** was synthesized from **III** (150 mg, 0.67 mmol) and lipoic acid (210 mg, 1.02 mmol) according to the procedure described for **RM11**, and purified by gravity column. Elution with petroleum ether/toluene/ $CH_2Cl_2$ /MeOH/ aqueous 30% ammonia (6:1:1.5:1.5:0.01) afforded **MR112** as a solid foam ceroso; yield 30%,  $^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.27 (t,  $J = 8.2$  Hz, 1H); 6.78-6.98 (m, 3H), 5.99 (br t, 1H); 4.09 (t,  $J = 6.0$  Hz, 2H); 3.21-3.62 (m, 5H), 3.05-3.19 (m, 3H), 2.40-2.53 (m, 1H), 2.32 (s, 6H), 2.22 (t,  $J = 7.2$  Hz, 2H); 1.81-2.0 (m, 3H), 1.65-1.73 (m, 4H), 1.47 (d,  $J = 6.6$  Hz, 3H); MS (ESI $^+$ )  $m/z$  411 (M+H) $^+$ . Calculated Anal. for  $C_{21}H_{34}N_2O_2S_2$ : C, 61.42; H, 8.35; N, 6.82. Found: C, 61.62; H, 8.36, N, 6.80.

<sup>i</sup> Present document

<sup>ii</sup> Chen, Yaun-Shek and Lawton, Richard G. An efficient synthetic route to 2-(1,2-dithiolan-3-yl)acetic acid. Trisnorlipoic acid and amide derivatives. *Tetrahedron Letters* **1997**, *38*, 5785-5788.

<sup>iii</sup> Patent UNIBO-LLG.

<sup>iv</sup> **I** was synthesized according to the procedure described for the corresponding (R,S)-3-[[1-di-( $^2H_3$ ) methylamino]ethyl]phenol in: Ciszewska, Grazyna; Pfefferkorn, Heidi; Tang, Y.S.; Jones, Lawrence; Tarapata, Richard; Sunay, Ustun B. Synthesis of tritium, deuterium, and carbon-14 labeled (s)-n-ethyl-n-methyl-3-[1-(dimethylamino)ethyl]carbamic acid, phenyl ester, (l)-2,3-dihydroxybutanedioic acid salt (SDZ ENA 713 hta), an investigational drug for the treatment of Alzheimer's disease. *Journal of Labelled Compounds & Radiopharmaceuticals* **1997**, *39*, 651-668.

## Figure and Scheme Legends

**Figure 1.** Design strategy for compounds 1-8.

**Figure 2.** Steady-state inhibition of AChE hydrolysis of acetylthiocholine (ACTh) by 7. Lineweaver-Burk reciprocal plots of initial velocity and substrate concentrations are reported. Reciprocal plots of initial velocity in the absence of inhibitor gave an estimate of  $k_{app}$  for acetylthiocholine of  $170 \pm 15 \mu\text{M}$  (four experiments). Lines were derived from a weighted least-squares analysis of the data points.

**Figure 3.** Effects of compounds on cell viability in neuronal cells. The cell viability was determined by the MTT assay (as described in the experimental section), after 24 h of incubation with various concentrations of LA (solid circle), 7 (open circle) and 15 (open triangle). The results were expressed as percentage of control cells. Values are reported as mean  $\pm$  SD of three independent experiments.

**Scheme 1.** Synthesis of compounds 1-8.




Figure 1




Figure 2





Figure 3



Scheme 1